

**APPENDIX A: Common and Scientific Names of Fishes Found in the  
Saint Mary - Belly River Recovery Unit**

northern pike; <i>Esox lucius</i> . . . . .	Native
white sucker; <i>Catostomus commersoni</i> . . . . .	Native
longnose sucker; <i>C. catostomus</i> . . . . .	Native
mountain sucker; <i>C. platyrhynchus</i> . . . . .	*Native
northern redbelly dace; <i>Phoxinus eos</i> . . . . .	*Native
pearl dace; <i>Margariscus margarita</i> . . . . .	Native
lake chub; <i>Couesius plumbeus</i> . . . . .	Native
longnose dace; <i>Rhinichthys cataractae</i> . . . . .	Native
fathead minnow; <i>Pimephales promelas</i> . . . . .	Native
kokanee; <i>Oncorhynchus nerka</i> . . . . .	Introduced
rainbow trout; <i>O. mykiss</i> . . . . .	Introduced
westslope cutthroat trout; <i>O. clarki lewisi</i> . . . . .	Native
brown trout; <i>Salmo trutta</i> . . . . .	*Introduced
brook trout; <i>Salvelinus fontinalis</i> . . . . .	Introduced
bull trout; <i>S. confluentus</i> . . . . .	Native
lake trout; <i>S. namaycush</i> . . . . .	Native
Arctic grayling; <i>Thymallus arcticus</i> . . . . .	Introduced
mountain whitefish; <i>Prosopium williamsoni</i> . . . . .	Native
lake whitefish; <i>Coregonus clupeaformis</i> . . . . .	Native
trout-perch; <i>Percopsis omiscomaycus</i> . . . . .	Native
burbot; <i>Lota lota</i> . . . . .	Native
brook stickleback; <i>Culaea inconstans</i> . . . . .	**Native
mottled sculpin; <i>Cottus bairdi</i> . . . . .	***Native
spoonhead sculpin; <i>C. ricei</i> . . . . .	Native

\* Not confirmed in U.S. (Montana) waters, but present in portions of the Saint Mary and/or Belly River drainages in Alberta (Nelson and Paetz, 1992).

\*\* Not present in Montana waters, limited to distribution in one tributary of the lower Saint Mary River drainage in Alberta (Nelson and Paetz, 1992).

\*\*\* In Alberta, the same fish is considered to be a variant of the shorthead sculpin *Cottus confusus*. More work is needed to resolve taxonomic questions.